



**awakenindiamovement** @awakenindiamov

Apr 17, 2023 · 70 tweets · [awakenindiamov/status/1648043305647603714](https://twitter.com/awakenindiamov/status/1648043305647603714)

DPT / DTP / TDap / DTaP thread



DTP, and DTP with OPV were both associated with increase in infant all cause mortality.  
(Mogensen et al.)

As a library, the National Library of Medicine (NLM) provides access to scientific literature.

Inclusion in an NLM database does not imply endorsement of, or agreement with, the contents by NLM or the National Institutes of Health.

**PubMed®**

**The Introduction of Diphtheria-Tetanus-Pertussis and Oral Polio Vacc...**  
DTP was associated with increased mortality; OPV may modify the effect of DTP.  
<https://pubmed.ncbi.nlm.nih.gov/28188123/>

> EBioMedicine. 2017 Mar;17:192-198. doi: 10.1016/j.ebiom.2017.01.041. Epub 2017 Feb 1.

## The Introduction of Diphtheria-Tetanus-Pertussis and Oral Polio Vaccine Among Young Infants in an Urban African Community: A Natural Experiment

Søren Wengel Mogensen <sup>1</sup>, Andreas Andersen <sup>2</sup>, Amabelia Rodrigues <sup>1</sup>, Christine S Benn <sup>3</sup>, Peter Aaby <sup>4</sup>

Affiliations + expand

PMID: 28188123 PMCID: [PMC5360569](https://pubmed.ncbi.nlm.nih.gov/28188123/) DOI: [10.1016/j.ebiom.2017.01.041](https://doi.org/10.1016/j.ebiom.2017.01.041)

[Free PMC article](#)

## Abstract


**Background:** We examined the introduction of diphtheria-tetanus-pertussis (DTP) and oral polio vaccine (OPV) in an urban community in Guinea-Bissau in the early 1980s.

**Methods:** The child population had been followed with 3-monthly nutritional weighing sessions since 1978. From June 1981 DTP and OPV were offered from 3months of age at these sessions. Due to the 3-monthly intervals between sessions, the children were allocated by birthday in a 'natural experiment' to receive vaccinations early or late between 3 and 5months of age. We included children who were <6months of age when vaccinations started and children born until the end of December 1983. We compared mortality between 3 and 5months of age of DTP-vaccinated and not-yet-DTP-vaccinated children in Cox proportional hazard models.

**Results:** Among 3-5-month-old children, having received DTP ( $\pm$ OPV) was associated with a mortality hazard ratio (HR) of 5.00 (95% CI 1.53-16.3) compared with not-yet-DTP-vaccinated children. Differences in background factors did not explain the effect. The negative effect was particularly strong for children who had received DTP-only and no OPV (HR=10.0 (2.61-38.6)). All-cause infant mortality after 3months of age increased after the introduction of these vaccines (HR=2.12 (1.07-4.19)).

**Conclusion:** DTP was associated with increased mortality; OPV may modify the effect of DTP.

Another study that observed that DTP only vaccines were associated with increased mortality in children 9-35 months of age,



**Findings:** The study cohort included 1491 children born in Bandim from December 1980 to December 1983. From 3 to 35 months of age, with censoring for MV, children vaccinated with DTP and/or OPV had higher mortality than both unvaccinated children (HR = 1.66; 95% CI, 1.03-2.69) and OPV-only vaccinated children (HR = 2.81; 95% CI, 1.02-7.69); DTP-only vaccinated children had higher mortality than OPV-only vaccinated children (HR = 3.38; 95% CI, 1.15--9.93). In the age group of 3-8 months, before MV is administered, DTP-only vaccination was associated with a higher mortality than DTP with OPV (HR = 3.38; 95% CI, 1.59-7.20). Between 9 and 35 months of age, when MV is given, DTP-vaccinated and MV-unvaccinated children had higher mortality (HR = 2.76; 95% CI, 1.36-5.59) than children who had received MV after DTP, and among children who received DTP with MV or after MV, DTP-only vaccination was associated with a higher mortality than DTP with OPV (HR = 6.25; 95% CI, 2.55-15.37).

"Receipt of DPT.... was associated with a possible increase in all cause mortality on average"...



#### Association of BCG, DTP, and measles containing vaccines with child...

Evidence suggests that receipt of BCG and MCV reduce overall mortality by more than would be expected through their effects on the diseases they prevent, and receipt of DTP may be associated with an ...


<https://pubmed.ncbi.nlm.nih.gov/27737834/>

Japan: DPT was temporarily suspended in 1974/75 following 2 deaths post vaccination. In 1981, DPT was replaced with DTaP, considered to be a "safer alternative"

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3824286/#!po=26.9231>

Fig. 4

#### History of pertussis vaccine and surveillance of pertussis



[Open in a separate window](#)

West Germany discontinued pertussis vaccine in 1974.

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2653494/>

Sweden discontinued the whole cell wP pertussis vaccine in 1979, and only included in the "safer" acellular pertussis vaccine in 1996




## Folkhälsomyndigheten

### Sidan kan inte hittas

Den sida du söker kan tyvärr inte hittas, men vi vill gärna hjälpa dig vidare. Använd i första hand sökfältet här ovanför för att söka det du letar efter.

<https://www.folkhalsomyndigheten.se/contentassets/cd49fff196f44e6a8db234ffb9da8b...>



"By the late 1990s, most high-income countries had switched to DTaP"

India - however continues to administer the whole cell pertussis component as part of it's DPT & pentavalent jabs, despite it's discontinuation in "first world" countries decades ago.



### Pertussis: a tale of two vaccines

Discover the world's best science and medicine | Nature.com

<https://www.nature.com/articles/d42859-020-00013-8>

Despite its known risks that led to its discontinuation in many countries, wholecell pertussis is a component of the 5 in 1 pentavalent vaccines, mortality from which, was found to be likely much higher than the DPT vaccines per one study.

awakenindiamovement  
@awakenindiamov · Follow

Replying to @awakenindiamov

" "This study demonstrated an increase in reports of sudden unexplained deaths within 72 hours of administering pentavalent vaccine compared to DTP vaccine." (Pulliyel, Jacob, et al.)

[journals.lww.com/mjdy/\\_layouts/...](journals.lww.com/mjdy/_layouts/...)

: AEFI deaths

	DTP vaccine	PV
Total vaccinated	44,936,653	24,803,770
Total deaths	217	237
Death rate per million vaccinated*	4.8	9.6

\* $P<0.0001$ . DTP: Diphtheria-tetanus-pertussis, PV: Pentavalent vaccine

11:05 PM · Mar 29, 2023

11 See the latest COVID-19 information on X

Read 2 replies

The havoc caused by the pentavalent vaccines in India is covered in the thread below.

awakenindiamovement  
@awakenindiamov · Follow

Replying to @awakenindiamov


Dr. Pulliyel estimated 1 in every 7,000 babies vaccinated with pentavalent  die.  
Isn't this unacceptable?

Did vaccines result in a decline in mortality from diphtheria & pertussis?


Example 1: Australia


Whooping cough death rate declined by 95% prior to introduction of vaccines  
<https://avn.org.au/information/vaccines-and-diseases/dtp/meryl/>

## The history of Whooping Cough in Australia





US: Mortality from whooping cough declined by over 90% prior to widespread use of vaccines






England & Wales - whooping cough deaths declined by ~99% prior to widespread mass immunization






France: Diphteria deaths were already on the decline prior to the introduction of vaccination. Oddly enough, the death rate increased after the introduction of vaccination.

[ia801707.us.archive.org/2/items/sincla](http://ia801707.us.archive.org/2/items/sincla)




Sweden: Diphteria deaths were already on the decline prior to the introduction of vaccination.

[https://ia801707.us.archive.org/2/items/sinclair-ian-vaccination-the-hidden-facts\\_202012/Sinclair%20Ian%20-%20Vaccination%20The%20hidden%20facts.pdf](https://ia801707.us.archive.org/2/items/sinclair-ian-vaccination-the-hidden-facts_202012/Sinclair%20Ian%20-%20Vaccination%20The%20hidden%20facts.pdf)



USA: Diphteria deaths were already on the decline prior to the introduction of vaccination.

[https://ia801707.us.archive.org/2/items/sinclair-ian-vaccination-the-hidden-facts\\_202012/Sinclair%20Ian%20-%20Vaccination%20The%20hidden%20facts.pdf](https://ia801707.us.archive.org/2/items/sinclair-ian-vaccination-the-hidden-facts_202012/Sinclair%20Ian%20-%20Vaccination%20The%20hidden%20facts.pdf)




Denmark: Diphteria deaths were already on the decline prior to the introduction of vaccination.

Death rate actually increased after the introduction of vaccines.

[https://ia801707.us.archive.org/2/items/sinclair-jan-vaccination-the-hidden-facts\\_202012/Sinclair%20Jan%20-%20Vaccination%20The%20hidden%20facts.pdf](https://ia801707.us.archive.org/2/items/sinclair-jan-vaccination-the-hidden-facts_202012/Sinclair%20Jan%20-%20Vaccination%20The%20hidden%20facts.pdf)


### Diphtheria: Denmark



England & Wales: Diphteria deaths were already on the decline prior to the introduction of vaccination.

[https://ia801707.us.archive.org/2/items/sinclair-ian-vaccination-the-hidden-facts\\_202012/Sinclair%20Ian%20-%20Vaccination%20The%20hidden%20facts.pdf](https://ia801707.us.archive.org/2/items/sinclair-ian-vaccination-the-hidden-facts_202012/Sinclair%20Ian%20-%20Vaccination%20The%20hidden%20facts.pdf)


## Diphtheria: England & Wales



Australia: Diphtheria deaths were already on the decline prior to the introduction of vaccination.

[https://ia801707.us.archive.org/2/items/sinclair-ian-vaccination-the-hidden-facts\\_202012/Sinclair%20Ian%20-%20Vaccination%20The%20hidden%20facts.pdf](https://ia801707.us.archive.org/2/items/sinclair-ian-vaccination-the-hidden-facts_202012/Sinclair%20Ian%20-%20Vaccination%20The%20hidden%20facts.pdf)

## Diphtheria: Australia



"In Norway diphtheria had virtually disappeared by 1939 when only 18 cases per million were recorded."

[https://ia801707.us.archive.org/2/items/sinclair-ian-vaccination-the-hidden-facts\\_202012/Sinclair%20Ian%20-%20Vaccination%20The%20hidden%20facts.pdf](https://ia801707.us.archive.org/2/items/sinclair-ian-vaccination-the-hidden-facts_202012/Sinclair%20Ian%20-%20Vaccination%20The%20hidden%20facts.pdf)

### Diphtheria

In Australia, diphtheria deaths for the period 1911-1915 were 3,677; 1921-1925 were 2,565; 1926-1930 were 1990; 1931-1935 were 2,083. Diphtheria vaccination commenced around 1932-1935 by which time a major reduction in the death numbers had already occurred. Dr Lancaster, referring to the decline in diphtheria, says:

*"... when the decline in mortality from diphtheria is compared with the decline in mortality rates from other childhood infections, it is seen that its relative decline has been no better than those of measles or pertussis (whooping cough) for which there was no specific treatment or prophylaxis up to the end of the period considered here".*

In England in 1860 diphtheria claimed annually over 1,000 deaths per million children, yet by 1870 this figure had fallen to around 400 deaths per million, even before the diphtheria germ had been isolated. By 1940 when diphtheria vaccination commenced, the annual death rate was down to less than 300 per year. From his book, Beyond The Magic Bullet, Bernard Dixon states:

*"Immunization against diphtheria, introduced on a large scale around 1940, appears to have had a dramatic effect on the incidence of the disease. The number of cases in Britain fell by between fifty and sixty thousand each year, until 1955, since when there have been only sporadic outbreaks. However, if we take a longer time scale, over the past century, and alter the criteria, we see a different picture. Diphtheria deaths in children went down continuously from 1300 per year in 1860, to under 300 per year in 1940, with a particularly*

5

### *Did Vaccines Really Save Us?*

*large drop around 1900, the year when antitoxin was first used. Yet the steepest decline was between 1865 and 1875 - before the diphtheria bacillus had even been isolated".*

Throughout Europe and America, diphtheria commenced its decline well before the introduction of diphtheria antitoxin, let alone vaccination. In Denmark, Sweden and Norway, deaths from diphtheria declined rapidly without vaccination. In Norway diphtheria had virtually disappeared by 1939 when only 18 cases per million were recorded.

Several instances of brain damage & death of healthy infants were reported in many countries per an article by Professor Gordon Stewart in 1980

[https://ia801707.us.archive.org/2/items/sinclair-ian-vaccination-the-hidden-facts\\_202012/Sinclair%20Ian%20-%20Vaccination%20The%20hidden%20facts.pdf](https://ia801707.us.archive.org/2/items/sinclair-ian-vaccination-the-hidden-facts_202012/Sinclair%20Ian%20-%20Vaccination%20The%20hidden%20facts.pdf)

### **Whooping Cough (Pertussis) Vaccine**

The vaccine against whooping cough is combined with vaccines for diphtheria and tetanus and is known as the DPT or triple antigen vaccine. Professor Gordon Stewart, in an article on whooping cough (Here's Health, March 1980) comments on the history of this vaccine in Britain.

*"Introduced in 1957, this vaccine had been administered to 70 per cent of infants by 1960 and over 70 per cent of all children by 1969.*

*"The national programme was monitored from 1957 - 1968 by the Public Health Laboratory Service. In 1969 they reported that the vaccines were 'not very effective' in that they had failed to control outbreaks or to protect fully-vaccinated children from infection. During this time the proportion of children vaccinated rose to 80 per cent or more and it is a matter of record that whooping cough continued to decline in prevalence and severity. But, equally, it is firmly on record not only that whooping cough occurred in fully-vaccinated children, but also that severe adverse reactions to the vaccine were causing problems and concern.*

*"If reference be made to events at the time of the earlier trials of pertussis vaccine when given alone (ie not as part of triple vaccine) in the USA and UK, it becomes clear that the inclusion of pertussis vaccine makes triple vaccine much more likely to be followed by adverse reactions involving the heart and nervous system. Such reactions include shock, collapse, convulsions and screaming fits, all of which had been recorded in some of the children who received pertussis vaccine alone in the earlier trials. Such signs were extremely infrequent or altogether absent in the earlier usage of the other two components of triple vaccine.*

*"More light was thrown on this problem when Professor W Ehrengut in Hamburg, and Dr John Wilson with colleagues at the Hospital for Sick Children, Great Ormond Street, London, reported independently that signs of severe brain damage began to appear in some children soon after adverse reactions to triple vaccine. At about the same time, a number of reports appeared in the Press from different parts of the UK about children who were previously well but had become mentally retarded or paralysed soon after receiving triple vaccine. The Government, on the advice of its advisory committees, responded to these reports by re-affirming the efficacy*

*and safety of pertussis vaccine and by insisting that this component be retained in triple vaccine. They insisted also that a high level of vaccination among children of all ages must be maintained if epidemics were to be averted.*

*"At that time in 1974, vaccination levels generally were about 80 per cent, seldom below 70 per cent and often above 90 per cent. The last outbreak of whooping cough had been in 1970-71 and, as epidemics are currently liable to occur every three to four years, another epidemic was expected and did in fact occur in 1974-75. This provided an opportunity for reviewing the efficacy of pertussis vaccine. It soon became apparent that protection was again incomplete and at best temporary, in that in all reports published at that time, a considerable proportion (30-50 per cent) of cases occurred in fully-vaccinated children.*

*"Meanwhile, reports about brain damage continued to circulate, leading to debates between experts and in Parliament about the safety of the vaccine. The main advisory committee (The Joint Committee on Immunization and Vaccination) stuck firmly to its view (first expressed in 1964) that the vaccine was safe as well as effective and that brain damage, if it occurred at all, was excessively rare, affecting no more than 1:300,000 infants vaccinated. They did, however, emphasise the need for caution, and recommend that the vaccine be withheld from children who showed signs of disorder in the nervous system, or had a family history of same, or who reacted badly to a first or second injection. There was by this time considerable doubt in many quarters, to which the government responded by setting up, through the Committee on the Safety of Medicines, a special expert panel to review the suspected toxicity of the vaccine. They also introduced in 1978 a scheme for compensation of parents of vaccine-damaged children.*

*"Between 1974 and 1978 acceptance of pertussis vaccine had been falling. Health authorities were offering a double vaccine (diphtheria plus tetanus) instead of a triple vaccine and this, together with poliomyelitis vaccine, was proving itself to be acceptable and unquestionably safe. But the Government's advisers were predicting a disastrous epidemic of whooping cough in the unprotected population. On the three to four year cycle, the next epidemic was due to begin in 1977, and it has to be acknowledged that notifications of whooping cough, which began to increase, then continued through 1978 and 1979 and amounted in total to what appeared to be the biggest epidemic since 1967. The mortality rate, however, was the lowest ever, and there was no doubt that the general pattern of previous epidemics was being followed, in that a high proportion of cases were observed among fully-vaccinated children. For the first time, there was a sharp difference in reports from different parts of the country. Some observers reported a low or even zero incidence in vaccinated children, while others found little difference between the vaccinated and unvaccinated.*

*"Internationally, the situation was equally confusing. In some countries like the USA and Canada, pertussis vaccine was used intensively and it was claimed that whooping cough was a disappearing disease. Nevertheless, in both of these countries outbreaks had been reported since 1974 in which (as in the UK) 30-50 per cent of cases were fully-vaccinated. In West Germany, largely as a result of Professor Ehrengut's work on toxicity, pertussis vaccine had been under suspicion for years and had been abandoned in Hamburg without any increase in incidence or mortality from*

*whooping cough. Similar decreases, without extensive use of vaccine, had occurred in Egypt and Italy.*

*"There is no doubt in my mind that in the UK alone some hundreds, if not thousands, of well infants have suffered irreparable brain damage needlessly and that their lives and those of their parents have been wrecked in consequence.*

*"There are also, to my certain knowledge, a number of deaths after vaccination in the UK and the USA which await explanation. I see no use or justification for this kind of medical policy, and I think that the use of pertussis vaccine should be discontinued until, by better research or a better vaccine, these doubts are resolved".*

## Observations from US

[https://ia801707.us.archive.org/2/items/sinclair-ian-vaccination-the-hidden-facts\\_202012/Sinclair%20Ian%20-%20Vaccination%20The%2ohidden%2ofacts.pdf](https://ia801707.us.archive.org/2/items/sinclair-ian-vaccination-the-hidden-facts_202012/Sinclair%20Ian%20-%20Vaccination%20The%2ohidden%2ofacts.pdf)

According to the Morbidity and Mortality Weekly Report (MMWR) July 5, 1985, a pertussis outbreak occurred in Washington, USA between January 1 to October 1, 1984 involving 162 cases. 69 of these cases occurred in children between 3 months old and 6 years. The report states: "*Of the 69 patients 3 months through 83 months (6 years) of age with known immunization status 34 (49%) were appropriately immunized for their ages with diphtheria and tetanus toxoids and pertussis vaccine*".

On September 2nd 1978, NBC News, Florida, USA, made the following announcement: "*The Atlantic Centre for Disease Control has asked doctors to stop using vaccines for diphtheria, tetanus and whooping cough because a number of children have been getting bad reactions*". Such "bad reactions" may include Sudden Infant Death Syndrome, commonly known as SIDS. According to Dr Alan Hinman, director of the Centre for Disease Control, Atlanta, USA: "*Since the CDC instituted its monitoring system in 1978, we have received reports of 44 deaths occurring within four weeks of DTP immunization. Thirty-two of the deaths were SIDS*". Leon Chaitow (Vaccination And Immunization) points to a study

19

### Vaccines - How Safe and Effective?

undertaken in 1979, at the University of California, Los Angeles (UCLA), under the sponsorship of the Food and Drug Administration, and which has been confirmed by other studies, indicates that in the USA approximately 1000 infants die annually as a direct result of DPT vaccination, and these are classified as SIDS deaths.

Dr William Torch of the University of Nevada, Reno, USA, has undertaken studies of SIDS cases. In one study, Dr Torch found that two thirds of 103 children who died of SIDS had been immunized with the DPT vaccine in the three weeks prior to their deaths. Many died within one day of vaccination. In 1982, Dr Torch, a noted pediatric neurologist said that the DPT vaccine "*may be a generally unrecognized cause of SIDS*".

From her book, I Had No Say, Sister Joyce Lubke writes:

*"When immunizations were given commencing at 3 months old, the peak of cot deaths was from 3-4 months. We are now told that the peak is 2-3 months, and this has happened since the immunizations commenced at 2 months. I feel there is some connection between cot deaths and immunization".*

In Health Report, Volume 6, No.12, December 1986, Dr H Buttram and J Hoffman tell of a study conducted by the Department of Paediatrics, University of California School of Medicine on 145 SIDS victims. Of this number, 53 had received DPT immunization shortly before their deaths. Of these 53, 27 died within a month of being vaccinated, 17 within a week and 6 within 24 hours.

From his book, How To Raise A Healthy Child In Spite Of Your Doctor, Dr Robert Mendelsohn writes:

*"My suspicion, which is shared by others in my profession, is that the nearly 10,000 SIDS deaths that occur in the US each year are related to one or more of the vaccines that are routinely given to children. The pertussis vaccine is the most likely villain, but it could also be one or more of the others".*

In their well researched 470 page book, DPT - A Shot in the Dark, co-authors H Coulter and B Fisher list potential side-effects and reactions to the DPT vaccine. They include skin reactions, fever, vomiting and diarrhoea, screaming and persistent crying, collapse, convulsions, infantile spasms, inflammation of the brain, blood disorders, diabetes, hypoglycaemia and SIDS.

In the USA, 1984, Edward Brandt Jnr, Assistant Secretary for Health, stated in a congressional testimony, that each year, the DPT vaccine will be associated with an estimated:

#### Vaccines - How Safe and Effective?

- 150 cases of brain inflammation or injuries, 50 with permanent damage;
- 9,000 cases of convulsion;
- 9,000 cases of collapse - a shock like state in which a child becomes limp, pale and unresponsive;
- 17,000 cases of unusual high-pitched screaming;
- 25,000 cases of fever of at least 105 degrees;
- 450,000 cases of inconsolable crying lasting from one to more than 20 hours.

In 1985 in the USA, an ABC Television research team, known as 20/20 uncovered massive amounts of documented evidence revealing the disastrous effects of DPT vaccine. 20/20 said that much of this information had been concealed by the drug companies, and that much of it was known by both Government and medical authorities who failed to take any action. 20/20 counted in excess of 2,500 cases of serious reactions, including brain damage, and over 60 deaths, all linked to the whooping cough vaccine. Evidence on the dangers of this vaccine went as far back as 1948. 20/20 found that government officials, doctors and vaccine manufacturers had held high level meetings on the dangers of this vaccine, without ever providing appropriate warning to the public!

“ Doctor Archie Kalokerinos has also observed the link between SIDS and immunization, noting that a number of apparently healthy aboriginal children, upon being vaccinated, would go into shock and die”

[https://ia801707.us.archive.org/2/items/sinclair-ian-vaccination-the-hidden-facts\\_202012/Sinclair%20Ian%20-%20Vaccination%20The%20hidden%20facts.pdf](https://ia801707.us.archive.org/2/items/sinclair-ian-vaccination-the-hidden-facts_202012/Sinclair%20Ian%20-%20Vaccination%20The%20hidden%20facts.pdf)

Doctor Archie Kalokerinos has also observed the link between SIDS and immunization, noting that a number of apparently healthy aboriginal children, upon being vaccinated, would go into shock and die. Speaking at a Natural Health Convention in Stanwell Tops, NSW, May 24th 1987, he had this to say about the whooping cough vaccine:

*“The worst vaccine of all is the whooping cough vaccine ... it is responsible for a lot of deaths and for a lot of infants suffering irreversible brain damage. In susceptible infants, it knocks their immune systems about, leading to irreparable brain damage, or severe attacks or even deaths from diseases like pneumonia or gastro-enteritis and so on”.*

US: “cost of a single DPT  had risen by 1,000% from.. 1982 .. to..1987. The manufacturers .. were putting aside \$8 per shot to cover legal costs & damages they were paying out to parents of brain damaged children & children who had died after immunization”

[https://ia801707.us.archive.org/2/items/sinclair-ian-vaccination-the-hidden-facts\\_202012/Sinclair%20Ian%20-%20Vaccination%20The%20hidden%20facts.pdf](https://ia801707.us.archive.org/2/items/sinclair-ian-vaccination-the-hidden-facts_202012/Sinclair%20Ian%20-%20Vaccination%20The%20hidden%20facts.pdf)

Besides the whole-cell pertussis component, another ingredient of the DPT vaccines, that was discontinued in pediatric vaccines in places like US & Canada, is 'Thiomersal', or ethylmercury, a known neurotoxin

[https://www.seruminstitute.com/product\\_ind\\_tripleantigen.php](https://www.seruminstitute.com/product_ind_tripleantigen.php)

<https://www.bharatbiotech.com/images/comvac3/comvac3-pres-info.pdf>

A booster dose can be given at the age of 15-18 months

## COMPOSITION

Each single 0.5 ml dose contains

Diphtheria Toxoid  $\leq$  25 Lf ( $\geq$  30 IU)

Tetanus Toxoid  $\geq$  5 Lf ( $\geq$  40 IU)

B. Pertussis  $\leq$  16 OU ( $\geq$  4 IU)

Adsorbed on Aluminium Phosphate, Al<sup>+++</sup>  $\leq$  1.25 mg

Preservative: 0.005% Thiomersal

## Composition

Each dose of 0.5 ml contains:

Diphtheria Toxoid	20 - 25 Lf
Tetanus Toxoid	5.0 - 7.5 Lf
B. Pertussis whole cell inactivated	15 OU - 20 OU
Aluminium Phosphate gel as Aluminium (Al <sup>+++</sup> )	0.3 mg
Thiomersal I.P. (as Preservative)	0.025 mg to 0.050 mg

There are studies that argue that the controversial ingredient - Thiomersal (mercury) is associated with neurodevelopmental disorders





Gyan Chakshu  
@gyan\_chakshu · Follow



Replying to @gyan\_chakshu  
[deeprootsathome.com/buried-evidenc...](http://deeprootsathome.com/buried-evidenc...)

6.

### Infants who received vaccines containing mercury developed speech disorders, sleep disorders and autism

*"This analysis suggests that high exposure to ethylmercury from thimerosal-containing vaccines in the first month of life increases the risk of subsequent development of neurologic development impairment."*

Verstraeten T, Davies R, et al. **Increased risk of developmental neurologic impairment after high exposure to thimerosal-containing vaccine in first month of life.** *Proceedings of the Epidemic Intelligence Service Annual Conference*, vol. 49 (Centers for Disease Control and Prevention; Atlanta, GA, USA, April 2000).

- This study was designed to determine whether infants who are exposed to ethylmercury from thimerosal-containing vaccines are at increased risk of degenerative and developmental neurologic disorders and renal disorders before the age of six.
- This study was conducted by the CDC using the Vaccine Safety Datalink (VSD) containing vaccination and demographic data on over 400,000 infants.
- The risk of developing a neurologic development disorder was nearly twice as high (RR = 1.8) in infants who received the highest cumulative exposure to ethylmercury (> 25 mcg) from thimerosal-containing vaccines at 1 month of age when compared to infants who were unexposed to mercury.
- One-month-old infants with the highest cumulative exposure to ethylmercury also had twice the risk of developing a speech disorder, 5 times the risk of developing a non-organic sleep disorder, and were 7.6 times more likely to develop autism when compared to infants who were unexposed to mercury from thimerosal-containing vaccines.

12:46 AM · Mar 29, 2022



28    Reply    Copy link

[Read 1 reply](#)

6.

## Infants who received vaccines containing mercury developed speech disorders, sleep disorders and autism

*"This analysis suggests that high exposure to ethylmercury from thimerosal-containing vaccines in the first month of life increases the risk of subsequent development of neurologic development impairment."*

Verstraeten T, Davies R, et al. **Increased risk of developmental neurologic impairment after high exposure to thimerosal-containing vaccine in first month of life.** *Proceedings of the Epidemic Intelligence Service Annual Conference*, vol. 49 (Centers for Disease Control and Prevention; Atlanta, GA, USA, April 2000).

- This study was designed to determine whether infants who are exposed to ethylmercury from thimerosal-containing vaccines are at increased risk of degenerative and developmental neurologic disorders and renal disorders before the age of six.
- This study was conducted by the CDC using the Vaccine Safety Datalink (VSD) containing vaccination and demographic data on over 400,000 infants.
- The risk of developing a neurologic development disorder was nearly twice as high (RR = 1.8) in infants who received the highest cumulative exposure to ethylmercury (> 25 mcg) from thimerosal-containing vaccines at 1 month of age when compared to infants who were unexposed to mercury.
- One-month-old infants with the highest cumulative exposure to ethylmercury also had twice the risk of developing a speech disorder, 5 times the risk of developing a non-organic sleep disorder, and were 7.6 times more likely to develop autism when compared to infants who were unexposed to mercury from thimerosal-containing vaccines.
- Premature babies were excluded from this study.
- There was no increased risk for neurologic degenerative and renal disorders.
- This study was never published.

Here are some media reports

May 2013 Daman: 1 child died, 3 were hospitalized following administration of DPT vaccine



May 2014, Agra, UP: A child died after DPT and measles vaccinations



January 2019, Mungoali MP, A 3 month old died following vaccination with DPT, 5 babies were hospitalized



October 2019: Mahasamund Chhattisgarh: A child was hospitalized in serious condition following DPT vaccination



July 2021, Bhadrak Odisha: A 2 month old baby died following OPV, DPT and Hepatitis vaccination.




2016, Hyderabad, Telangana: Child died after DPT and polio vaccination.



December 2019: Abdasa, Gujarat: Child died following DPT and measles rubella vaccination



May 2017, Muzzafarpur Bihar: 2 children died and 12 hospitalized following tetanus and Japanese encephalitis vaccination



August 2022, Sednawa Madhya Pradesh: 18 children hospitalized following tetanus and diphtheria vaccinations




July 2021, Banaskantha Gujarat: 3 children hospitalized following diphtheria vaccination



Baby died following measles and tetanus vaccination.

Date: January 2020

Place: Giridih, Jharkhand



5 year old child died following diphtheria and pertussis vaccinations.

February 2021, Keeratpur, Uttar Pradesh




8 month old baby died following DPT vaccination.

December 2021, Hazaribagh Jharkhand



One child died and 17 hospitalized following DPT vaccination.

August 2020, Bagha, Bihar



Child died following DPT vaccination.

Place: Darbhanga, Bihar

Date: December, 2014



2022, October, Kolhapur Maharashtra - 1.5 year old infant died following vaccination

Per immunization schedule, Measles rubella, DPT, Japanese encephalitis and oral polio are administered at this age



Two 3 month old babies died following and another 2 hospitalized following vaccination for BCG and DPT

Date: November 2018

Place: Parbhani, Maharashtra



Shameful @MoHFW\_INDIA @myogiadityanath :

!!FORCED VACCINATION!!

150 children forcibly vaccinated with TD / DPT vaccine. 50 children sick.

Place: Aligarh, Uttar Pradesh

Date: October, 2022



अलीगढ़ के स्कूल में 150 बच्चों को जबरन दी गई वैक्सीन डोज, 50 से अधिक बच्चों की बिगड़ी हालत  
Aligarh Primary School: अलीगढ़ के सरकारी विद्यालय में पढ़ने वाले लगभग 150 बच्चों को वैक्सीन की डोज लगा दी गई, वैक्सीन की डोज लगने के बाद लगभग 50 से अधिक बच्चों की तबीयत बिगड़ गई।

<https://www.abplive.com/states/up-uk/aligarh-news-vaccine-dose-was-forcibly-given-to...>



**Aligarh, Uttar Pradesh, 150 children forcibly vaccinated with DT and D...**

October 2022, Aligarh, Uttar Pradesh, 150 children forcibly vaccinated with DT and DPT vaccines, 50 fell sick <https://www.abplive.com/states/up-uk/aligarh-news-vaccine-dose-was-forcibly-given-to-150-...>

<https://rumble.com/v2my7qi-aligarh-uttar-pradesh-150-children-forcibly-vaccinated-wit...>

**!!FORCED VACCINATION!!**

15 adolescent girls hospitalized following tetanus vaccination. Apparently, parents were not informed, no consent was taken.

@MoHFW\_INDIA @BhagwantMann @narendramodi @PMOIndia @ArvindKejriwal



<https://www.youtube.com/embed/5ZEhPJFANw8>



**Big News: पंजाब के School में लगे Injection से 10 और Students की बिगड़ी हालत, दे...**  
इंजेक्शन लगाने से 10 और छात्राओं की हालत बिगड़ी

<https://punjab.punjabkesari.in/punjab/news/10-students-experience-side-effects-after-t...>



Date: May 11th, 2023

Place: Ludhiana, Punjab

 **awakenindiamovement**  
@awakenindiamov · [Follow](#) X

Replying to @awakenindiamov  
**!! FORCED VACCINATION !!**


15 adolescent girls hospitalized following tetanus vaccination. Apparently, parents were not informed, no consent was taken.

@MoHFW\_INDIA @BhagwantMann @narendramodi  
@PMOIndia @ArvindKejriwal

[youtube.com/watch?v=5ZEhPJ...](https://youtube.com/watch?v=5ZEhPJ...)

[punjab.punjabkesari.in/punjab/news/10...](https://punjab.punjabkesari.in/punjab/news/10...)

**Figure 1. Polio cases skyrocketed after diphtheria and pertussis vaccines were introduced**



11 year old girl died following vaccination in school.

Per immunization schedule tetanus and diphteria vaccinations are administered at around this age

Date: August, 2022

Place: Dahod, Gujarat

@MoHFW\_INDIA @mansukhmandviya @narendramodi

Dharmik vala  
@dharmik\_vala · Follow

X

દાહોદના વડબારા ગામમાં ધનુરની રસી આપતા મોત વડબારા પ્રાથમીક શાળામાં મુકવામાં આવી હતી રસી 11 વર્ષિય બાળકીના મોતથી પરીવારમાં શોક ધનુરની રસીના કારણે મોત થયાનો આદ્દેપ

#dahod #Gujarat



Watch on X



On the occasion of "Mamta Diwas", children were vaxxed

One such child, Varshaben, a 4 year old girl, died . DPT vacs r given at 16-24 m & 5-6 years of age per imm. sched.


Date: June 29th, 2023

Place: Khambaliya, Gujarat

<https://khabargujarat.com/innocent-girl-from-khambaliya-dies-after-injection-of-vaccine/>



<https://www.youtube.com/embed/DsdvazfB-aA>



January 2021, Khurja, Bulandshahar, UP: 18 month old infant died following vaccination.

Per immunization schedule, Measles rubella, DPT, Japanese encephalitis and oral polio are administered at this stage

[https://public.app/video/sp\\_ybwl38k4ihsho](https://public.app/video/sp_ybwl38k4ihsho)



5 year old boy hospitalized, admitted to critical care following vaccination in school

Per immunization schedule, DPT vaccine is administered at this stage

Date: November, 2019

Place: Cuddalore, Tamil Nadu

@MoHFW\_INDIA @mansukhmandviya @narendramodi



<https://www.youtube.com/embed/LjLbLkUldto>



July 2016, Salem, Tamil Nadu, 3 children hospitalized following mass vaccination campaign in school

5-6 years of age - Per imm; sched., DPT administered at this stage

10 years of age - Per imm. sched, -Tetanus and diphteria administered at this stage



<https://www.youtube.com/embed/pvLVI7z4mIQ>



18 children hospitalized following Tetanus Diphtheria vaccination in school.

Date: August 2022

Place: Sendhwa, Madhya Pradesh

Zee MP-Chhattisgarh ✅  
@ZeeMPCG · Follow

टीकाकरण के बाद बच्चों की बिगड़ी तबीयत, सेंधवा में लगाया गया था  
टिटनेस और डिप्टीरिया का टीका

#MadhyaPradesh

@Pooja\_Sachdeva\_

For More Updates : [bit.ly/3AaaapX](https://bit.ly/3AaaapX)

Watch on X

4:46 AM · Aug 20, 2022

1 8 Reply Copy link

[Read more on X](#)



Toddler hospitalized due to infection in arm following vaccination

Date: September 18th 2023

Place: Hathras Uttar Pradesh

@MoHFW\_INDIA @myogiadityanath <https://hamarahathras.com/hathras-365878-2/>  
[dailymotion.com/video/x8o6985](https://dailymotion.com/video/x8o6985)



Shameful @MoHFW\_INDIA @myogiadityanath @mansukhmandviya @narendramodi



Awaken India Movement @awakenindiamvmt · Follow



!! 80 children sick following tetanus and DPT vaccination in government school !!

The parents have alleged that no parental consent was taken prior to vaccinating the children.

@PMOIndia @MoHFW\_INDIA @myogiadityanath  
@mansukhmandviya - why is this gross violation of  
parental... Show more





Awaken India Movement

@awakenindiamvmt · Follow



!! The carnage of ongoing Tetanus Diphtheria (TD) vaccination campaign continues. !!

Ayushi, a 7 year old girl has died following vaccination. 4 other children sick.

Officials blaming mid day meal

@MoHFW\_INDIA @myogiadityanath @narendramodi  
@mansukhmandviya Where's the... Show more



Awaken India Movement

@awakenindiamvmt · Follow



Replies to @awakenindiamvmt

!! 2 year old toddler died following multiple vaccinations  
 !!

Date: December 21st, 2023

Place: Hubbali, Karnataka

Per immunization schedule, Measles rubella, DPT, Japanese encephalitis and oral polio are administered at this age

@CMofKarnataka - are you going to simply... Show more



Anuraj, a 1.5 year old infant died 11 hours after DPT vaccination

Date: December 23, 2023

Place: Paroo, Bihar



# हिन्दुस्तान

देवरियाकोठी में डेढ़ साल के मासूम की संदिग्ध हालत में मौत  
देवरिया थाना के विश्वनपुर सरैया गांव के जयपरकाश नगर में शुक्रवार देर रात एक डेढ़ साल के बच्चे की संदिग्ध हालत में मौत हो गयी। जिसकी पहचान रामपरवेश... Muzaffarpur News Hindustan

<https://www.livehindustan.com/bihar/muzaffarpur/story-death-of-one-and-a-half-year-old-child-in-vishwanapur-saraiya-village>



Bihar News: मुजफ्फरपुर के पारू में टीकाकरण के ग्यारह घंटे के अंदर बच्चे की मौत, महकमे में...  
बिहार के मुजफ्फरपुर में पारू पर्स्संड के विश्वनपुर सरैया पंचायत में एक डेढ़ वर्षीय बच्चे की जीवीटी टीकाकरण दिए जाने के ग्यारह घंटे के भीतर शुक्रवार की रात मौत हो गयी। उसे गांव के आंगनबाड़ी केंद्र पर दिन क...

<https://www.jagran.com/bihar/muzaffarpur-child-dies-within-eleven-hours-of-vaccination>



<https://www.youtube.com/embed/pBkrld-As8>



Kapil, a 6 year old boy died following vaccination

Date: August 2022

Place: Mandla, Madhya Pradesh

Vaccination was performed without consent of parents.

Per immunization schedule, DPT is administered at this age

मोहगांव की मानोट स्कूल का मामला; पिता बोले- बिना बताए टीका गया, अस्पताल पहुंचने से पहले ही ...  
मंडला: मंडला जिले की मोहगांव जनपद अंतर्गत ग्राम मानोट की प्राथमिक शाला में टीका लगाने के बाद एक 6 वर्षीय छात्र कपिल नंदा की मौत हो गई। छात्र के पिता का आरोप है कि उनके बेटे को स्कूल में बिना सूचना दिए...

<https://naisochlive.com/59133/>

[https://public.app/video/sp\\_oxcswzjgijgoi](https://public.app/video/sp_oxcswzjgijgoi)





**Awaken India Movement** [@awakenindiamvmt](#) · [Follow](#)



Replying to @awakenindiamvmt

18 month old baby died following DPT vaccination

Date: February 19th, 2024

Place: Kalaburagi, Karnataka

YTD: 14

Feb: 6

[m.dinamalar.com/detail.php?id=...](http://m.dinamalar.com/detail.php?id=...)

[etvbharat.com/kn/lstate/it-i...](http://etvbharat.com/kn/lstate/it-i...)

[vistaranews.com/karnataka/kala...](http://vistaranews.com/karnataka/kala...)

[facebook.com/gundurao.bhing...](http://facebook.com/gundurao.bhing...)

[facebook.com/watch/?ref=sea.....](http://facebook.com/watch/?ref=sea.....)



#Tetanus vaccine and fertility issues



Awaken India Movement

@awakenindiamvmt · [Follow](#)



Tetanus vaccine may be laced with anti-fertility drug.  
[pubmed.ncbi.nlm.nih.gov/12346214/](https://pubmed.ncbi.nlm.nih.gov/12346214/) [Show more](#)

> Vaccine Wkly. 1995;9:10.

## Tetanus vaccine may be laced with anti-fertility drug. International / developing countries

No authors listed

PMID: 12346214

### Abstract

**PIP:** A priest, president of Human Life International (HLI) based in Maryland, has asked Congress to investigate reports of women in some developing countries unknowingly receiving a tetanus vaccine laced with the anti-fertility drug human chorionic gonadotropin (hCG). If it is true, he wants Congress to publicly condemn the mass vaccinations and to cut off funding to UN agencies and other involved organizations. The natural hormone hCG is needed to maintain pregnancy. The hormone would produce antibodies against hCG to prevent pregnancy. In the fall of 1994, the Pro Life Committee of Mexico was suspicious of the protocols for the tetanus toxoid campaign because they excluded all males and children and called for multiple injections of the vaccine in only women of reproductive age. Yet, one injection provides protection for at least 10 years. The Committee had vials of the tetanus vaccine analyzed for hCG. It informed HLI about the tetanus toxoid vaccine. HLI then told its World Council members and HLI affiliates in more than 60 countries. Similar tetanus vaccines laced with hCG have been uncovered in the Philippines and in Nicaragua. In addition to the World Health Organization (WHO), other organizations involved in the development of an anti-fertility vaccine using hCG include the UN Population Fund, the UN Development Programme, the World Bank, the Population Council, the Rockefeller Foundation, the US National Institute of Child Health and Human Development, the All India Institute of Medical Sciences, and Uppsala, Helsinki, and Ohio State universities. The priest objects that, if indeed the purpose of the mass vaccinations is to prevent pregnancies, women are uninformed, unsuspecting, and unconsenting victims.

>Last edited 6:52 AM · Apr 27, 2024



66



[See the latest COVID-19 information on X](#)

[Read 4 replies](#)



Awaken India Movement

@awakenindiamvmt · [Follow](#)



!! SCHOOL CHILDREN VACCINATED WITHOUT PARENTAL CONSENT, 1 GIRL DIED !!

27 children were vaccinated with the TD vaccine, without parental consent in a school in Silchar, Assam. One of them, an 11 year old girl died 3 days after vaccination

Date: May 3rd, 2024

Place: Silchar,... [Show more](#)



awakenindiamovement  
@awakenindiamov · [Follow](#)



Thread on [#dtap](#)

● **Gyan Chakshu** @gyan\_chakshu


#DTap

Child suffered generalized epilepsy with febrile seizures plus ("GEFS+") following DTaP vaccination.

Court ordered over 32 million dollars in compensation (\$1 million lump sum + annuities totalling \$31.4 million)

[casetext.com/case/harris-v-...](http://casetext.com/case/harris-v-...)

[uscfc.uscourts.gov/sites/default/...](http://uscfc.uscourts.gov/sites/default/...)



8:43 AM · May 13, 2024



2



See the latest COVID-19 information on X

[Read more on X](#)

This is an article about the 11 babies who died in Tennessee between 1978-79 after receiving a certain batch of pertussis/DTaP vaccines. Four of them died within 24 hours.

<https://circleofmamas.com/health-news/the-tennessee-sids-cluster/>



Lauren Houston [Follow](#)



This is an article about the 11 babies who died in Tennessee between 1978-79 after receiving a certain batch of pertussis/DTaP vaccines. Four of them died within 24 hours. I find the story about Wyeth/Pfizer subsequently hiding "hot lots" incredibly disturbing.... [Show more](#)

7:36 PM · Mar 25, 2024



21



See the latest COVID-19 information on X

[Read 2 replies](#)



Awaken India Movement @awakenindiamvmt · [Follow](#)



!! 3 year old child died following vaccination !!



Date: April 18th, 2024

Place: Habra, West Bengal

16-24 months - Per immunization schedule, Measles rubella, DPT, Japanese encephalitis and oral polio are administered at this age

5-6 - Per immunization schedule, DPT... [Show more](#)



Awaken India Movement @awakenindiamvmt · [Follow](#)



✳️ 1.5 year old baby suffered seizures following DPT vaccination, died following treatment ✳️



Date: May 31st, 2024

Place: Patiala, Punjab

[youtube.com/watch?app=desk...](https://youtube.com/watch?app=desk...)

@Tleensthack  
@PriyankaPulla  
@RPrasad12  
@ShamikaRavi  
@IASassociation  
@themojostory  
@BDUTT  
@ndtv  
@aajtak... [Show more](#)





Awaken India Movement @awakenindiamvmt · [Follow](#)



\*\*\*\*7 schoolgirls hospitalized following TD vaccination in school\*\*\*\*

Date: July 27th, 2024

Place: Bulandshahar, Uttar Pradesh

#SaveTheChildren

[m.youtube.com/watch?v=B2S816...](https://m.youtube.com/watch?v=B2S816...)

[bhaskar.com/amp/local/utta...](https://bhaskar.com/amp/local/utta...)


[m.up.punjabkesari.in/uttar-pradesh/...](https://m.up.punjabkesari.in/uttar-pradesh/...)

[amarujala.com/amp/delhi-ncr/...](https://amarujala.com/amp/delhi-ncr/...)

@JPNadda

@MoHFW\_INDIA

@PMOIndia... [Show more](#)



Watch on X



Awaken India Movement @awakenindiamvmt · [Follow](#)



SHAMEFUL!! 6 schoolgirls hospitalized after all students were forcibly vaccinated 😢. TD vaccine was administered.

Do watch the video to listen to the testimony of the girls who were forcibly vaccinated. UNACCEPTABLE !!

Bureaucrats, politicians, medicos all sing praises of the... [Show more](#)



Watch on X

2013 study from Haryana which showed that there was significant excess deaths among girl kids after the #DTP vaccine.



As a library, the National Library of Medicine (NLM) provides access to scientific literature.  
Inclusion in an NLM database does not imply endorsement of, or agreement with, the contents by NLM or the National Institutes of Health.

PubMed®


**Non-specific sex-differential effect of DTP vaccination may partially ex...**  
This study supports the contention that DTP vaccination is partially responsible for higher mortality among girls in this study population.  
<https://pubmed.ncbi.nlm.nih.gov/24103109>

 **Neo**  
@neorevolt · [Follow](#)

X

2013 study from Haryana which showed that there was significant excess deaths among girl kids after the DTP vxxine.

There are similar studies from Africa and couple of other countries also.



**Non-specific sex-differential effect of DTP vaccination may partially explain the excess girl child mortality in Ballabgarh, India**

A Krishnan <sup>1</sup>, R Srivastava, P Dwivedi, N Ng, P Byass, C S Pandav  
Affiliations + expand  
PMID: 24103109 DOI: [10.1111/tmi.12192](https://doi.org/10.1111/tmi.12192)  
[Free article](#)

**Abstract**

**Objective:** To test the hypothesis that a gender differential exists in the effect on child mortality of BCG, DTP, measles vaccine as administered under programme conditions in Ballabgarh HDSS area.

**Methods:** All live births in 28 villages of Ballabgarh block in North India from 2006 to 2011 were followed until 31 December 2011 or 36 months of age whichever was earlier. The period of analysis was divided into four time periods based on eligibility for vaccines under the national immunisation schedule (BCG for tuberculosis, primary and booster doses of diphtheria-tetanus-pertussis and measles). Cox proportional hazards regression was used to assess the association between sex and risk of mortality by vaccination status using age as the timescale in survival analysis and adjusting for wealth index, access to health care, the presence of a health facility in the village, parental education,

6:01 AM · Sep 3, 2024

 35   Copy link

[Read more on X](#)



## Non-specific sex-differential effect of DTP vaccination may partially explain the excess girl child mortality in Ballabgarh, India

A Krishnan <sup>1</sup>, R Srivastava, P Dwivedi, N Ng, P Byass, C S Pandav

Affiliations + expand

PMID: 24103109 DOI: 10.1111/tmi.12192

[Free article](#)

### Abstract

**Objective:** To test the hypothesis that a gender differential exists in the effect on child mortality of BCG, DTP, measles vaccine as administered under programme conditions in Ballabgarh HDSS area.

**Methods:** All live births in 28 villages of Ballabgarh block in North India from 2006 to 2011 were followed until 31 December 2011 or 36 months of age whichever was earlier. The period of analysis was divided into four time periods based on eligibility for vaccines under the national immunisation schedule (BCG for tuberculosis, primary and booster doses of diphtheria-tetanus-pertussis and measles). Cox proportional hazards regression was used to assess the association between sex and risk of mortality by vaccination status using age as the timescale in survival analysis and adjusting for wealth index, access to health care, the presence of a health facility in the village, parental education, type of family, birth order of the child and year of birth.

**Results:** 702 deaths (332 boys and 370 girls) occurred among 12,142 children in the cohort in the 3 years of follow-up giving a cumulative mortality rate of 57.5 per 1000 live births with 35% excess girl child mortality. Age at vaccination for the four vaccines did not differ by sex. There was significant excess mortality among girls after immunisation with DTP, for both primary (HR 1.65; 95% CI:1.17-2.32) and DTPb (2.21; 1.24-3.93) vaccinations. No significant excess mortality among girls was noted after exposure to BCG 1.06 (0.67-1.67) or measles 1.34 (0.85-2.12) vaccine.

**Conclusion:** This study supports the contention that DTP vaccination is partially responsible for higher mortality among girls in this study population.

DTP vaccine study from Africa shows it was associated with 5-fold higher deaths in the vaccinated kids than kids who haven't taken the vaccine



Neo

@neorevolt · Follow

X

DTP vxxine study from Africa shows it was associated with 5-fold higher deaths in the vxxed kids than kids who haven't taken the vxxine. And even those kids who didn't die, no prospective study has shown any beneficial survival effects of the vxxine.

The screenshot shows a research paper from the journal EBioMedicine. The title is "The Introduction of Diphtheria-Tetanus-Pertussis and Oral Polio Vaccine Among Young Infants in an Urban African Community: A Natural Experiment". The authors listed are Søren Wengel Mogensen, Andreas Andersen, Amabelia Rodrigues, Christine S Benn, and Peter Aaby. A key sentence in the abstract is underlined: "DTP was associated with 5-fold higher mortality than being unvaccinated. No prospective study has shown beneficial survival effects of DTP. Unfortunately, DTP is the most widely used vaccine, and the proportion who receives DTP3 is used globally as an indicator of the performance of national vaccination programs." Below the abstract, there is a paragraph about the concern over routine vaccinations and their effect on all-cause mortality. At the bottom of the screenshot, there are social media interaction icons (heart, reply, copy link) and a link to "Read more on X".

EBioMedicine  
journal homepage: www.ebiomedicine.com

Research Paper  
The Introduction of Diphtheria-Tetanus-Pertussis and Oral Polio Vaccine Among Young Infants in an Urban African Community: A Natural Experiment  
Søren Wengel Mogensen <sup>a,1</sup>, Andreas Andersen <sup>b,1</sup>, Amabelia Rodrigues <sup>a</sup>, Christine S Benn <sup>b,c</sup>, Peter Aaby <sup>a,b,\*</sup>

DTP was associated with 5-fold higher mortality than being unvaccinated. No prospective study has shown beneficial survival effects of DTP. Unfortunately, DTP is the most widely used vaccine, and the proportion who receives DTP3 is used globally as an indicator of the performance of national vaccination programs.

It should be of concern that the effect of routine vaccinations on all-cause mortality was not tested in randomized trials. All currently available evidence suggests that DTP vaccine may kill more children from other causes than it saves from diphtheria, tetanus or pertussis. Though a vaccine protects children against the target disease it may simultaneously increase susceptibility to unrelated infections.

The recently published SAGE review called for randomized trials of DTP (Higgins et al., 2014). However, at the same time the IVIR-AC committee to which SAGE delegated the follow-up studies of the NSEs of vaccines has indicated that it will not be possible to examine the effect of DTP in an unbiased way. If that decision by IVIR-AC remains unchallenged, the present study may remain the closest we will ever come to a RCT of the NSEs of DTP.

6:10 AM · Sep 3, 2024

1

9

Reply

Copy link

Read more on X

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5360569/>



## Research Paper

## The Introduction of Diphtheria-Tetanus-Pertussis and Oral Polio Vaccine Among Young Infants in an Urban African Community: A Natural Experiment



Søren Wengel Mogensen <sup>a,1</sup>, Andreas Andersen <sup>b,1</sup>, Amabelia Rodrigues <sup>a</sup>, Christine S Benn <sup>b,c</sup>, Peter Aaby <sup>a,b,\*</sup>

DTP was associated with 5-fold higher mortality than being unvaccinated. No prospective study has shown beneficial survival effects of DTP. Unfortunately, DTP is the most widely used vaccine, and the proportion who receives DTP3 is used globally as an indicator of the performance of national vaccination programs.

It should be of concern that the effect of routine vaccinations on all-cause mortality was not tested in randomized trials. All currently available evidence suggests that DTP vaccine may kill more children from other causes than it saves from diphtheria, tetanus or pertussis. Though a vaccine protects children against the target disease it may simultaneously increase susceptibility to unrelated infections.

The recently published SAGE review called for randomized trials of DTP (Higgins et al., 2014). However, at the same time the IVIR-AC committee to which SAGE delegated the follow-up studies of the NSEs of vaccines has indicated that it will not be possible to examine the effect of DTP in an unbiased way. If that decision by IVIR-AC remains unchallenged, the present study may remain the closest we will ever come to a RCT of the NSEs of DTP.

...